

For Research Use Only

# FcZero-rAb™ PE-Cyanine7 Anti-Mouse CD11b (M1/70) Rabbit IgG Recombinant Antibody

Catalog Number: PY7-FcA65672



## Basic Information

Catalog Number:

PY7-FcA65672

Size:

100ug , 200 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC156991

GeneID (NCBI):

16409

Full Name:

integrin alpha M

Calculated MW:

127 kDa

Purification Method:

Protein A purification

CloneNo.:

M1/70

Recommended Dilutions:

FC: 0.25 ug per 10<sup>6</sup> cells in a 100 µl suspension

Excitation/Emission maxima wavelengths:

450-500 nm / 778 nm

## Applications

Tested Applications:

FC

Species Specificity:

human, mouse

Positive Controls:

FC : mouse bone marrow cells,

## Background Information

Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated  $\alpha$  and  $\beta$  subunits (PMID: 9779984). CD11b, also known as Integrin alpha M or CR3A, belongs to the integrin alpha chain family. CD11b forms an  $\alpha/\beta$  heterodimer with CD18 (integrin  $\beta$ 2). CD11b/CD18 is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complement-coated particles and pathogens (PMID: 9558116; 20008295). CD11b/CD18 is a receptor for the complement protein fragment iC3b, and is also a receptor for fibrinogen, factor X and ICAM1 (PMID: 2971974; 15485828).

## Storage

Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA, pH7.3


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10<sup>6</sup> mouse bone marrow cells were surface stained with 0.25 ug PE-Cyanine7 Anti-Mouse CD11b (M1/70) Rabbit IgG RecAb (PY7-FcA65672, Clone: M1/70) or 0.25 ug PE-Cyanine7 Rabbit IgG Isotype Control RecAb (PY7-FcA98136, Clone: 240953C9). Cells were incubated with FcZero Monocytes Block prior to staining. Cells were not fixed.