

For Research Use Only.
Not For Use In Diagnostics.

Multi-rAb™ Polymer HRP-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L)



Catalog Number:RGAR011

Information

Catalog Number:

RGAR011

Host:

Goat

Applications:

IHC

Reactivity:

Rabbit

Physical State:

Liquid

Conjugation:

Polymer HRP

Clonality:

Multiclonal recombinant

Recommended Dilutions

Ready to use

Safety Notes

This product is for research use only, not for diagnostic or therapeutic use.

Storage

Storage:

Store at 2-8°C. Stable for one year after shipment.

Storage Buffer:

TBST+10 mg/mL BSA+1 mg/mL gelatin + 0.1% proclin300

Aliquoting is unnecessary for -20°C storage

Purity

Protein G purification


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB
(1-888-478-4522)
(toll free in USA),
or 1(312) 455-8498
(outside USA)


E : proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 80001-1-RR (TDP43 recombinant antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-Rabbit Recombinant secondary antibody RGAR011 was used for detection.




Immunohistochemical analysis of paraffin-embedded mouse colon tissue slide using 80008-1-RR (smooth muscle actin recombinant antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection.




Immunohistochemical analysis of paraffin-embedded rat colon tissue slide using 80008-1-RR (smooth muscle actin recombinant antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection.




Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 12715-1-AP (Dystrophin polyclonal antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection.




Immunohistochemical analysis of paraffin-embedded mouse colon tissue slide using 20874-1-AP (E-cadherin polyclonal antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection.




Immunohistochemical analysis of paraffin-embedded mouse skin tissue slide using 20874-1-AP (E-cadherin polyclonal antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection.




Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 80165-1-RR (TMEM173/STING antibody) at dilution of 1:2000. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using commercial IVD grade Ki67 antibody. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using other vendor's NF-kappa B p65 rabbit recombinant antibody. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection.




Immunohistochemical analysis of paraffin-embedded human lymphoma tissue slide using other vendor's BCL2 rabbit recombinant antibody. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection.




Immunohistochemical analysis of paraffin-embedded rat ovary tissue slide using 12255-1-AP (RCAS1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.




Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue slide using 12255-1-AP (RCAS1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.




Immunohistochemical analysis of paraffin-embedded rat skeletal muscle tissue slide using 12715-1-AP (Dystrophin antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer. Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.




Immunohistochemical analysis of paraffin-embedded mouse tongue tissue slide using 12715-1-AP (Dystrophin antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.




Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 12715-1-AP (Dystrophin antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.




Immunohistochemical analysis of paraffin-embedded mouse skin tissue slide using 20874-1-AP (E-cadherin antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.




Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 20874-1-AP (E-cadherin antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.




Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 80001-1-RR (TDP-43 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 80001-1-RR (TDP-43 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.




Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 80001-1-RR (TDP-43 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.




Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 81668-1-RR (F4/80 antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.




Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 81668-1-RR (F4/80 antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection. The staining was performed on Celnovte's CNT360-M1 autostainer.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using other vendor's ER rabbit recombinant antibody. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection.



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 21146-1-AP (LARS antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection.



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 21146-1-AP (LARS antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Multi-rAb™ Polymer HRP-Goat anti-rabbit Recombinant secondary antibody RGAR011 was used for detection.