For Research Use Only

DYKDDDDK Fab-Trap® Agarose, Kit for Immunoprecipitation



www.ptglab.com

Catalog Number: ffak

8 Publications

Basic Information

Catalog Number:

Applications: IP, CoIP, ChIP, RIP, Protein purification

Agarose beads; bead size: ~ 90 µm (cross-linked 4 % agarose beads)

Host: Mouse

Type: Fab-fragment

Class: Recombinant

Description

The ChromoTek DYKDDDDK Fab-Trap® Agarose Kit consists of an anti-DYKDDDDK-tag Fab fragment, which is covalently bound to agarose beads. DYKDDDDK Fab-Trap® Agarose Kit is used to immunoprecipitate 1x and 3xDYKDDDDK-tagged fusion proteins from cell extracts of various organisms like mammals, plants, bacteria, yeast, insects etc

Binding capacity

12.5 - 15 µg of recombinant 3xDYKDDDDK-tagged protein (~30 kDa) per 25µL bead slurry

Specificity/Target

The DYKDDDDK Fab-Trap® Agarose specifically binds to 1x and 3xDYKDDDDK-tag sequence (1x and 3xFlag®-tag). It binds to same epitope as Sigma's Anti-FLAG® Antibody and is compatible with N-, C-terminal or internal tagging.

Elution buffer


3xDYKDDDDK-peptide SDS sample buffer 0.2 M glycine pH 2.5

Storage


Shipped at ambient temperature. Upon receipt store at 4°C. Stable for one year. Do not freeze!

Storage Buffer: 1x PBS, Preservative: 0.09 % sodium azide


Selected Validation Data



DYKDDDDK Fab-Trap® was used for immunoprecipitation of Flag®-tagged protein from HEK293T cell lysate. During elution with 3xDYKDDDDK-peptide (fp-1) the enriched Flag® fusion protein is released from DYKDDDDK Fab-Trap™. Western blot was probed with DYKDDDDK tag Polyclonal antibody (Binds to FLAG® tag epitope) (Proteintech, 20543-1-AP) and Nano-Secondary® alpaca anti-human IgG/anti-rabbit IgG, recombinant VHH, Alexa Fluor® 488 [CTK0101, CTK0102] (srbAF488-1). L: Prestained protein marker (Proteintech, PL00001), I: Input, FT: Flow-Through, E: Elution.



DYKDDDDK Fab-Trap® was used for immunoprecipitation of Flag®-tagged protein from HEK293T cell lysate. During elution with 2x SDS-sample buffer the enriched Flag® fusion protein and Fab-fragment are released from DYKDDDDK Fab-Trap™. Western blot was probed with DYKDDDDK tag Polyclonal antibody (Binds to FLAG® tag epitope) (Proteintech, 20543-1-AP) and Nano-Secondary® alpaca anti-human IgG/anti-rabbit IgG, recombinant VHH, Alexa Fluor® 488 [CTK0101, CTK0102] (srbAF488-1). L: Prestained protein marker (Proteintech, PL00001), I: Input, FT: Flow-Through, B: Bound.



DYKDDDDK Fab-Trap® was used for immunoprecipitation of Flag®-tagged protein from HEK293T cell lysate. During elution with 3xDYKDDDDK-peptide (fp-1) the enriched Flag® fusion protein is released from DYKDDDDK Fab-Trap™. Western blot was probed with DYKDDDDK tag Polyclonal antibody (Binds to FLAG® tag epitope) (Proteintech, 20543-1-AP) and Nano-Secondary® alpaca anti-human IgG/anti-rabbit IgG, recombinant VHH, Alexa Fluor® 488 [CTK0101, CTK0102] (srbAF488-1). L' Prestained protein marker (Proteintech, PL00001), I: Input, FT: Flow-Through, E: Elution.