

# Nano-Secondary® alpaca anti-mouse IgG2b, recombinant VHH, CoraLite® Plus 488 [CTK0105, CTK0106]

Product code: smsG2bCL488-1

### **Properties**

**Description** Monovalent, recombinant secondary single domain antibody to mouse IgG2b:

Mixture of 2 alpaca monoclonal Nanobodies, Fc-specific, CoraLite® Plus 488

conjugated

Product type Nano-Secondary® Reagent, secondary Nanobody (VHH)

**Format** Alpaca single domain antibody, monovalent

**Host** Alpaca-derived, recombinantly produced in bacteria

**Target/Specificity** Fc-fragment of mouse IgG2b

**Cross-reactivity** No cross-reactivity to goat, guinea pig, human, macaque, rabbit, rat, and

sheep serum and to mouse IgG1, IgG2a, IgG2c, IgG3, and IgM

**Immunogen** Purified mouse IgG2b

**Clonality** Biclonal: mixture of 2 monoclonal Nanobodies

Clones CTK0105 (VHH0275), CTK106 (VHH0288)

**Affinity (**Kd**)** CTK0105: KD = 5 nM, CTK106: KD = 0.2 nM

**Conjugate** CoraLite® Plus 488

**Excitation / Emission** Excitation max: 490 nm, Emission max: 525 nm

**Degree of labeling (DOL)** 2 fluorophores per Nanobody

Synonyms Alpaca single domain antibody, VHH, Nanobody, binding domain of single

domain antibody, Nano-antibody

**Validation** Application validated for immunofluorescence and western blotting.

Determination of cross-reactivity, sequence, affinity, melting point, and degree

of labeling (DOL).

**Purity** Recombinantly expressed and purified

**Form** Buffered aqueous solution

**Concentration** 0.5 mg/mL

**Storage buffer** 10 mM HEPES pH 7.0, 500 mM NaCl, 5 mM EDTA

Preservative: 0.09 % sodium azide, safety datasheet (SDS): sodium azide

**Storage instructions** Shipped at ambient temperature. Store at +4°C short term or -20°C long term.

Stable for 1 year at -20°C.

**Size** 10  $\mu$ L; 100  $\mu$ L

**RRID** AB\_2941314



## Nano-Secondary® alpaca anti-mouse IgG2b, recombinant VHH, CoraLite® Plus 488 [CTK0105, CTK0106]

Product code: smsG2bCL488-1

**Applications** Immunofluorescence: recommended dilution range of 1:500-1:1000.


Western blot: recommended dilution range of 1:500-1:1000.

The optimal dilution should be determined by the user. A titration range is

recommended.


## **Tested applications**

#### *Immunofluorescence*



Immunofluorescence analysis of HeLa cells stained with mouse IgG2b anti-Tubulin beta antibody and Nano-Secondary® alpaca antimouse IgG2b, recombinant VHH, CoraLite® Plus 488 (smsG2bCL488-1, green). Nuclei were stained with DAPI (blue). Images were recorded at the Core Facility Bioimaging at the Biomedical Center, LMU Munich

#### Western blot



HEK-293 cell lysates were subjected to SDS-PAGE followed by multiplex western blot analysis with 3 mouse primary antibodies including anti-Lamin B1 (66095-1-lg), anti-HNRNPA (67445-1-lg), and anti-Tom20 (66777-1-lg). Primary antibodies were detected using 3 mouse lgG subclass-specific nano-secondary reagents including Nano-Secondary® alpaca anti-mouse lgG1, recombinant VHH, CoraLite® Plus 647 (smsG1CL647-1, blue), Nano-Secondary® alpaca anti-mouse lgG2a, recombinant VHH, CoraLite® Plus 750 (smsG2aCL750-1, gray), and Nano-Secondary® alpaca anti-mouse lgG2b, recombinant VHH, CoraLite® Plus 488 (smsG2bCL488-1, green).

Only for research applications, not for diagnostic or therapeutic use.

ChromoTek is a registered trademark of ChromoTek GmbH, part of Proteintech Group. CoraLite is a registered trademark of Proteintech Group. Nanobody is a registered trademark of Ablynx, a Sanofi company. Other suppliers' products may be trademarks or registered trademarks of the corresponding supplier each. Statements on other suppliers' products are given according to our best knowledge.