For Research Use Only

Nano-Secondary® anti-mouse IgG1, recombinant VHH, CoraLite® Plus 555 [CTK0103, CTK0104]



www.ptglab.com

Catalog Number: smsG1CL555-1

Catalog Number: smsG1CL555-1 **Basic Information**

Applications: IF, WB

Alpaca-derived, recombinantly produced in bacteria

CoraLite® Plus 555

Type: Nanobody Class: Recombinant **RRID:** AB_2941311

Purification Method:

Recombinant expression, affinity purification

Description Nano-Secondary® anti-mouse IgG1, Fc-specific recombinant VHH is an anti-mouse IgG subclass specific antibody. This

secondary antibody product consists of Nanobodies that bind to mouse IgG1 with high affinity & specificity.

Species Reactivity Mouse IgG1 Fc-fragment

 $No\ cross \'- reactivit \'y\': goat, guinea\ pig,\ rabbit,\ rat,\ sheep,\ human,\ macaque\ serum\ proteins,\ mouse\ lgG2a,\ lgG2b,\ lgG2c,\ lgG3,\ lgMacapha lgMacapha$

Physical State Liquid

Suggested Dilution Immunofluorescence 1:500


Western blot 1:500

Affinity CTK0103: KD =0.13 nM, CTK0104: KD = 0.63nM


Storage Shipped at ambient temperature. Store at +4°C short term or -20°C long term. Stable for 1 year at -20°C.

Storage Buffer: 10 mM HEPES pH 7.0, 500 mM NaCl, 5 mM EDTA, Preservative: 0.09 % Sodium azide


Selected Validation Data



Immunofluorescence analysis of HeLa cells stained with mouse IgG1 anti-HSP60 antibody (66041-1-Ig) and Nano-Secondary® alpaca anti-mouse IgG1, recombinant VHH, Coralite® Plus 555 (smsG1CL555-1, orange). Nuclei were stained with DAPI (blue). Images were recorded at the Core Facility Bioimaging at the Biomedical Center, LMU Munich



Immunofluorescence analysis of HeLa cells costained with mouse IgG1 anti-HSP60 antibody (66041-1-Ig) and mouse IgG2b anti-lamin A/C antibody followed by Nano-Secondary® alpaca anti-mouse IgG1, recombinant VHH, Coralite® Plus 555 (smsG1CL555-1, yellow) and Nano-Secondary® alpaca anti-mouse IgG2b, recombinant VHH, Coralite® Plus 647 (smsG1CL647-1, magenta). Images were recorded at the Core Facility Bioimaging at the Biomedical Center, LMU Munich.



HEK-293 cell lysates were subjected to SDS-PAGE followed by multiplex western blot analysis with 3 mouse primary antibodies including anti-TDP43 (60019-1-Ig), anti-tubulin (66240-1-Ig), and anti-IMP3 (66247-1-Ig). Primary antibodies were detected using 3 mouse IgG subclass-specific nanosecondary reagents including Nano-Secondary® alpaca anti-mouse IgG1, recombinant VHH, Coralite® Plus 555 (smsG1CL555-1, red), Nano-Secondary® alpaca anti-mouse IgG2a, recombinant VHH, Coralite® Plus 647 (smsG2aCL647-1, blue), and Nano-Secondary® alpaca anti-mouse IgG2b, recombinant VHH, Coralite® Plus 750 (smsG2bCL750-1, white).