For Research Use Only

## Myc-Trap® 2.0 Agarose, Kit for **Immunoprecipitation**



www.ptglab.com

Catalog Number: yt2ak

**Description** 

**Catalog Number: Basic Information** 

**Applications:** IP, Co-IP

Conjugate: Agarose beads; ~90 um (cross-linked 4% agarose beads)

Type: Nanobody Class: Recombinant

**Host:** 

Alpaca

The ChromoTek Myc-Trap® Agarose, Kit for Immunoprecipitation consists of an anti-Myc NANOBODY®/VHH, which is coupled to agarose beads. It also contains lysis, wash, and elution buffers that can be used for the immunoprecipitation of Myc-fusion

proteins from cell extracts of various organisms.

Binds specifically to the Myc-tag (sequence EQKLISEEDL) at the N-terminus, C-terminus, or internal site of the fusion protein. Endogenous c-myc is NOT bound. Specificity/Target

**Elution buffer**  $2x\ SDS-sample\ buffer\ (L\"{a}mmli),\ 200\ mM\ glycine\ pH\ 2.5,\ 0.1\ mg/ml\ ChromoTek\ 2x\ Myc-peptide\ (2yp)\ in\ PBS\ pH\ 7.4$ 


Affinity (K<sub>D</sub>)

Storage


Shipped at ambient temperature. Upon receipt store at +4°C. Stable for one year. DO not freeze!

Storage Buffer: 20% ethanol

## **Selected Validation Data**



Immunoprecipitation of GFP-Myc fusion protein from HEK293T cells using Myc-Trap® 2.0 Agarose. IP was done using both untransfected (mock) and transfected (GFP-myc) cells. I: Input: F: Flow-through, B: Bound.



Comparison of pulldown efficacy between the Myc-Trap® 2.0 Agarose (left) and the original Myc-Trap Agarose (right). Both products were used to immunoprecipitate GFP-myc fusion proteins from untransfected (mock) and transfected (GFP-myc) HEK293T cells. The Myc-Trap 2.0 has higher affinity for myc-tagged proteins and is able to pulldown more GFP-Myc protein than the Myc-Trap.